

The Solar System

Magda Stavinschi, Beatriz García, Andrea Sosa

International Astronomical Union


Astronomical Institute of the Romanian Academy, Romania

ITeDA and National Technological University, Argentina

University of the Republic, Uruguay




Here is where stars are born




Messier 16, Pillars of creation.
Credit: Hubble Space Telescope

The Orion Nebula

and here



Hubble Space Telescope
Wide Field Planetary Camera 2



The planets in the past: the naked eyes visibles

Mercury
Venus



Visible at
sunset or
sunrise

Mars

Jupiter

Saturne

Planetary alignment,
May- 2002



The Solar System today

It is constituted by the Sun and all the bodies that revolve around it, under the action of gravity:

- 8 planets
- Hundred of natural satellites of planets
- Tenths of dwarf planets (between them Ceres, Pluto, Haumea, Makemake and Eris)
- An unknown number of minor bodies: asteroids, comets and transneptunians (debris of the processes of the planet formation).



Where is the Solar System?

It is in **Orion arm**, one of the **Milky Way** arms.

"Spiral Galaxy, similar to Milky Way"



The Milky Way has about 200,000 millions of stars and its diameters is about 100,000 l.y.

The Solar System is located at a distance of $\sim 25,000$ light years from the center of the Galaxy (\sim half the radius), and takes 250 millions years to complete a rotation around the center. The speed is 220 km/s (800.000 km/h)



Model of the Milky Way, from the infrared observations by Spitzer (2005); our Galaxy is a barred spiral.

Solar System formation

- According to the standard theory, about 4.6 billion years ago the solar system was formed from the gravitational contraction of an interstellar gas and dust cloud. The collapse of the cloud started from a strong perturbation (possibly a supernova burst), which caused the gravitational force to overcome the pressure of the gases.



- The conservation of the angular momentum caused the nebula to turn faster and faster, to flatten out, and to give rise to a protostar at its center, and to a protoplanetary disk of gas and dust around it.

Solar System formation


- In the protoplanetary disk condensed small solid nuclei (planetesimals), which then were accumulated by an accretion process to form the planets.
- The standard theory described above is accepted for having found, through high resolution radio images, protoplanetary systems around many young stars and due to the possibility of explaining the formation of planets within those systems.



Solar System Studies

The Sun concentrates more than 99.8% of the mass of the SS, while 98% of the angular momentum is found in the orbital movements of the planets.

Currently, the study of solar system bodies is done from Earth, but also through space telescopes, sending missions to space and even descending on its surface.





Our star: The Sun

- With an age of 4,600 million years, the Sun is approximately in the middle of its life cycle.
- Every second, in the Sun's core, 4 million tons of matter are converted into energy, generating a large number of neutrinos, positrons and radiation.





74% of the Sun is H, 25% is He, the rest are heaviest elements.

Sun structure




Energy production: fusion in the core.

The life of the stars depends on their masses




Life cycle of the Sun

Within 5,000 million years, the Sun will swell and become a red giant. Then it will expel the outer layers, creating a planetary nebula, and in the center there will be a small star called white dwarf, which will slowly cool down.



The Planets



XXVI IAU-AG Resolution, Praha, 2006:

In the SS, a “**planet**” It is a celestial body that:


- It is in orbit around the Sun.
- It has enough mass for its self-gravity (which is a central force) to impose itself on the cohesive forces of a rigid body so that it assumes a form in hydrostatic (quasi-spherical) equilibrium.
- It has cleared other objects the neighborhood along its orbit.

A body that meets only the first two criteria, and that is not a satellite, is classified as a “**dwarf planet**”.

A body that meets only the first criteria, and that is not a satellite, is called “small body (or **minor body**) of the SS.

The Solar System today


(bodies in size scale)



The limit of the Solar System



All planetary orbits lie within the Heliosphere, a region of space that contains magnetic fields and plasma ("wind") of solar origin.

The Heliopause is the limit of the Heliosphere, where the solar wind merges with the interstellar medium.



In 2012, the Voyager 1 space probe crossed the Heliopause at a heliocentric distance of more than 100 A.U.


Earth shown
for size comparison



Sun-Earth environment

The Interplanetary Medium

The Sun emits electromagnetic radiation and solar wind (a continuous flow of charged particles, plasma).



This dissipates at a speed of 1.5 million km / h, creating the heliosphere, a fine atmosphere that bathes the entire SS up to approx. 100 A.U., marking the heliopause.

The Earth's magnetic field protects the atmosphere from the solar wind and gives rise to the polar auroras (boreal and austral)



The heliosphere ensures a partial protection to the SS of the cosmic rays, protection that is stronger in the planets with magnetic field.

The “space weather” is monitoring 24 hs

SpaceWeather.com -- News and information about meteor showers, solar flares, auroras, and near-Earth asteroids - Mozilla Firefox

File Edit View History Bookmarks Tools Help

http://www.spaceweather.com/

Google cinturones de van allen Search Share Bookmarks Check Translate AutoFill cinturones

SpaceWeather.com -- News and info...

 **spaceweather.com**
News and information about the Sun-Earth environment

Subscribe to SpaceWeatherNews go!

AURORA ALERTS | **SUBMIT YOUR PHOTOS!** | **3D SUN** | **CONTACT US** | **SUBSCRIBE** | **FLYBYS** | **SCIENCE@NASA**

Current Conditions

Solar wind
speed: **347.4** km/sec
density: **1.1** protons/cm³
[explanation](#) | [more data](#)
Updated: Today at 0546 UT

X-ray Solar Flares
6-hr max: **B8** 0032 UT Mar29
24-hr: **B8** 0032 UT Mar29
[explanation](#) | [more data](#)
Updated: Today at: 0500 UT

Daily Sun: 28 Mar 11



What's up in space

Tuesday, Mar. 29, 2011

Metallic photos of the sun by renowned photographer Greg Piepol bring together the best of art and science. Buy one or a whole set. They make a stellar gift.



SOLAR RADIO STORM: Did you know sunspots can make noise? Consider the following: "Over the past few days, I have been recording a sustained solar radio storm at 180 MHz," reports amateur radio astronomer [Thomas Ashcraft](#) of New Mexico. "It consists of Type I radio bursts and sounds like ocean surf. [Here is an audio sample](#) from March 27th at 1930 UT. The sun seems to be entering a new phase of dynamism."

Radio emissions like these are caused by plasma instabilities in the sun's atmosphere above sunspots. With the sun becoming 'radio-active,' it's no coincidence that sunspots are emerging in abundance. Leading the way is behemoth active region AR1176, shown here in a photo taken yesterday by Larry Alvarez of Flower Mound, Texas:



archives

March

29

2011

space toys.com


Averted Imagination
ASTROPHOTOGRAPHY

The Planets

The 8 planets of our SS can be divided into:

- **4 Earth planets**, in the innermost region (Mercury, Venus, Earth and Mars). Rocky, with approximate densities between 4 and 5 g / cm³.
- **4 Giant planets**, in the outermost region, which in turn are divided into:
 - **Gaseous Giants**: Jupiter and Saturn. Richer in H and He, with a chemical composition similar to solar.
 - **Ice Giants**: Uranus and Neptune. Ice predominates with respect to gases. Its chemical composition differs a lot from solar.
- The giant planets are lighter than terrestrial ones, with densities between 0.7 g/cm³ (Saturn) and 2 g/cm³.

The giant planets had formed on time scales of the order of 10 million years (terrestrial planets did in about 100 million years).

They were not formed "in situ", there was a migration caused by the exchange of angular momentum between the giant planets in formation and the planetesimals that were swept to other regions of the SS or ejected from the SS


Earth



Earth-Moon system,
photographed by the
Galileo spacecraft,
on its way to Jupiter
(1998)





April 12, 1961
First circumnavigation
flight to Earth by
Yuri Gagarin



Mercury

The one closest to
the Sun, presents
an impact surface






The most important crater is "Caloris Basin" (1,500 km in diameter): the impact that originated it produced waves that broke the surface at the antipodes (photo).

The precession of perihelion of Mercury

The precession of Mercury's perihelion is faster than the predictions of Newton's classical celestial mechanics.

That advance of perihelion was predicted by Einstein's General Theory of Relativity.


It is due to the curvature of the space caused by the Sun. It was a definitive proof of that Theory.



Venus



Observed on
Earth with an
small telescope



Observed by the
Hubble Space
Telescope




VENERA (1976)

ВЕНЕРА-9 22.10.1975

ОБРАБОТКА ИППИ АН СССР 28.2.1976




Magellans (1990-1994)



With dimensions and geological structure similar to those of the Earth, it was visited by several missions

Venus and Uranus are the only planets with retrograde movement (they turn on themselves in the opposite direction to how they revolve around the Sun).



- **Venusian year = 224 Earth days**
- **Venusian day = 243 Earth days.**

The mixture of CO_2 and dense clouds of sulfur dioxide create the greatest greenhouse effect of the entire SS, with temperatures reaching 460°C , higher than that of Mercury.

The atmospheric pressure is 100 times the Earth's pressure, there are clouds and perhaps rain of sulfuric acid.

Venus Transit


When Venus passes between the Earth and the Sun, its shadow crosses the solar disk.

Because the inclination of the orbit of Venus occurs twice in 8 years, and the next takes more than a century (105.5 or 121.5 years).

In June 2004 and June 2012 the last ones took place. There will not be another until December 11, 2117



Mars



It has a fine atmosphere,
composed mainly of CO₂.
The atmospheric pressure is
one hundredth of the Earth's.






Primera imagen de
Marte, Viking I, 1976



First photo on Mars
surface Viking I, 1976

Source of inspiration for many science fiction authors ("extraterrestrial" = "Martian"), due to the famous "canali" observed by Giovanni Schiaparelli at the end of the 19th century: the term was translated into English as "canals" as if they were human constructions.



Its red color is due to the oxide of Fe (hematite), which is found in surface minerals





Cydonia – Viking I, 1976





Cydonia

Mars Global Surveyor 1998



Cydonia -Mars Express – Sep., 2006



There are traces that indicate that there was water on Mars.



The water may now be frozen in the subsoil.



As on Earth, there are stations on Mars because the axis of rotation is inclined with respect to the orbital plane, and because the planets move around the Sun keeping constant the inclination of the axis.



Martian Southern Pole

It has two ice caps, ice and CO₂ whose extension varies with the seasons.



Curiosity on Mars (2004-present): a successful history of science and technology: a microbiology laboratory



Insight: arrives on Mars on Nov 28, 2018

InSight (Interior Exploration using Seismic Investigations, Geodesy and Heat Transport,)




OBJECTIVE: to place a geophysical robot, equipped with high-tech instruments to study the interior, subsoil, heat transmission and movements of the Martian soil and analyze the early geological evolution of the planet.

Jupiter

A detailed view of Jupiter's atmosphere, showing the Great Red Spot as a large, reddish-orange oval storm. Other smaller, swirling cloud patterns in various shades of brown, tan, and white are visible across the planet's surface.

The most massive planet of the SS, has more than 60 moons. In 1610 Galileo observed for the first time 4 of them that he called "Mediceas". That same year Simon Marius baptized them as Io, Europa, Ganymede and Callisto.

Auroras, Photo by Hubble Telescope



Large red dot (a cyclone)



Probably has a small solid core, between 10 and 15 times the Earth's mass.




Anillos de Júpiter

Rings system

Saturn

The less dense planet of the SS.



It has more than 60 moons and some of them are between the rings, dynamically organizing the system, they are called "shepherd satellites"


System of Rings, formed by dust and very small pieces of ice.



Aurora in Saturn, photo by the Hubble Space Telescope




- Saturne has more than 60 satellites but 7 are large enough to take a spherical shape.
- Titan is the largest (larger than Mercury and Pluto) and the only one in the SS with a dense atmosphere.



Cassini-Huygens Mission



Huygens probe
descending on Titan
(artistic vision)







Sonda Huygens on Titan (first panoramic photo, 2004)

Titan: seas, rivers and lakes of methane





Last photo
on Titan
surface,
Huygens
probe




Uranus

Its axis of rotation is practically in its plane of translation




Rings system of Uranus





Uranus has at least 27 natural satellites.

The first two were discovered by William Herschel in 1787: Titania and Oberon.



The satellites of Uranus bear the names of the heroines of Shakespeare's plays





Urano • Julio 28, 1997

HST • NICMOS

PRC97-36a • November 20, 1997 • ST ScI OPO


E. Karkoschka (University of Arizona Lunar & Planetary Lab) and NASA

Neptune



Its color is due to the presence of methane in the atmosphere, which absorbs red and infrared light.

Neptune





It is believed that it has a solid core of silicates and iron, almost as large as Earth.

Above the core is a shell of ice, methane, H and a little He

It has several dark rings, with origin unknown.





**Clyde Tombaugh,
discovered Pluto
on February 18th,
1930.**




**Image of the
discovery. (1930)**

Pluto is too small to disturb Neptune's orbit long enough to betray its presence, however much Lowell has calculated to locate it. Clyde Tombaugh found Pluto (magnitude ~ 13.5) photographing in a systematic way the plane of the SS.




Pluto Faces
Hubble Space Telescope • ACS/HRC

Pluto System ■ February 15, 2006
Hubble Space Telescope ■ ACS/HRC



NASA, ESA, H. Weaver (JHU/APL), A. Stern (SwRI),
and the HST Pluto Companion Search Team

Pluto System, 2011-2012







Pluto and Charon
New Horizons, 2015

20 miles

NASA




**Overflight of Pluto
(July 14, 2015)
The faint atmosphere
of nitrogen is observed**



Eris Discovery

Planeta enano Eris y satélite Dysnomia. Agosto 30, 2006.


HST • ACS/HRC



Minor bodies of the Solar System

- **They are the remnants of the planetary accretion.**
- They comprise diverse populations of asteroids, comets and transneptunian objects.
- The asteroids are essentially rocky and metallic, while the comets are more fragile and porous objects, formed basically by ice (predominantly water) and dust particles.
- The vast majority of asteroids lie in a region between the orbits of Mars and Jupiter, known as the "Asteroid Main Belt."
- Transneptunian objects will contain significant amounts of ice, and are located in a region beyond the orbit of Neptune, known as the "Transneptunian Belt" (or Kuiper Belt, in recognition of one of the first to predict its existence).

Asteroids Main Belt




There are hundreds of thousands or millions, and the total mass would not exceed one thousandth of the Earth.

The size of the asteroids ranges from several hundred km to meters and fractions of m.

Ceres


Discovered in 1801 by Giuseppe Piazzi, it was considered a planet until 1850 when many other similar objects were found.

It is the largest body of the asteroid belt, and the only one of them cataloged in 2006 as a dwarf planet.




With a diameter of almost 1,000 km, it is large enough for its gravity to give it a spherical shape.

All other asteroids are considered small, irregular bodies, although some of them like Pallas and Vesta could be classified as dwarf planets if they are shown to reach hydrostatic equilibrium



Pallas



Reservoirs of minor bodies in the SS

The reservoirs are relatively stable regions, where objects can remain for times comparable to the SS age, until some perturbative force changes their orbit.

There are three large reservoirs in the SS:


- The **Main Asteroid Belt** Other populations would come from this region, such as the asteroids that approach Earth (known as NEAS by its acronym in English).
- The **Transneptunian Belt**. It is the region where short-period comets come from.
- The **Oort Cloud**. It has a spherical distribution and is formed by the frozen planetesimals swept out by giant planets during the formation of the SS. Thanks to perturbations due to the close passage of stars or giant molecular clouds, or to the galactic tides, the orbits of some of these objects can change deviating towards the interior of the SS, transforming into long-period comets.

Data at April 17, 2019.


Source: NASA/JPL (<https://ssd.jpl.nasa.gov>)

- **Total of known Asteroids: 798,130. Including:**
 - **Main belt: 705,913**
 - **Trojans of Jupiter: 7,236**
 - **Asteroids with inner Mars orbits: 3,573**
 - **NEAs: 19,996**
 - **Partial dangerous Asteroids (PHAs): 1,973**
- **Comets:**
 - **Elliptical: 420 long period (P>200 years) + 860 short period (P<200 years).**
 - **Parabolics: 1,837**
 - **Hyperbolic: 347 (extra-solar origin)**
- **Trans-neptunians (TNOs): 3,218**

Transneptunian belt and Oort cloud




Trans
neptunians



The Oort cloud
(comprising many
billions of comets)

The
biggest
ones are
dwarf
planets

Largest known trans-Neptunian objects (TNOs)



2000 km

Comets

- ❑ Are small bodies of a few km, made mainly of volatile materials (water ice, carbon dioxide, methane, ammonia, etc.) and dust particles.
- ❑ When they approach the Sun they can be visible.
- ❑ It is thought that H₂O on Earth could come from them.




West, 1976



Hale-Bopp, 1997

- In general comets have quite eccentric orbits. Those of long period have random inclinations and may have retrograde or direct orbits: those of short period have generally small inclinations and their orbits are direct.

- When approaching the Sun, the superficial ice of the comet is sublimated creating a coma or "hair", and the "tails": a dust tail formed by dust particles dragged by the gas, and an ionic tail formed by the atoms and ionized molecules that interact with the solar wind. The powder tail is curved, while the bluish ionic tail points straight and opposite the Sun




Halley: the most famous of the comets

It was named in honor of Edmond Halley, who predicted his approach to the Sun, applying the Law of Universal Gravitation and the calculation of disturbances. Halley did not see his prediction confirmed.



It returns every 76 years.




Distance: 3625 km
Time to encounter: 52 sec




In 1986 was the first comet visited by a probe: the Giotto. It photographed the nucleus.

Rosetta Mission: close encounter with the comet 67P/Churyumov-Gerasimenko


Philae descends on the comet on Nov 12, 2014



67P surface



Nucleus Activity (Sept. de 2014)



Camera OSIRIS/ESA

Other Planetary Systems

In 1995 the Swiss astronomers Michel Mayor and Didier Queloz announced the detection of an exoplanet orbiting 51 Pegasi.




□ This star and its planet were baptized as Helvetios and Dimidio in 2015, after a public vote promoted by the IAU.

1st photo of an extrasolar planet around a brown dwarf 2M1207. March 16, 2003


Exoplanet Discoveries Through the Years

As of May 10, 2016



Kepler's Planets by Size

As of May 10, 2016




Kepler (March 2009), is NASA's first mission to find potentially habitable planets, the size of Earth.

On May 10, 2016, he announced the largest exoplanet collection for which news is available.

Out of a total of about 5,000 candidates, more than 3,200 have been verified, and 2,325 of these were discovered by the Kepler telescope.

Since 2018, the NASA satellite "Transiting Exoplanet Survey" will use the same method as the Kepler telescope to monitor 200,000 nearby bright stars and search for planets, especially the size of Earth or greater (the super Earths).



How many stars have planets?

How many of those planets are habitable?

In how many developed some form of
life?

Questions that astronomy
seeks to answer

**Many Thanks
for your attention!**

